



*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*

Transforming matter and dark matter into antimatter

Zurab Berezhiani

University of L'Aquila and LNGS

GSSI Vanity Fair, 21 Feb. 2020





Contents

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 $n - n'$ and
UHECR

- 1 Introduction: Dark Matter from a Parallel World
- 2 Chapter I: *Neutrino - mirror neutrino mixings*
- 3 Chapter II: *neutron - mirror neutron mixing*
- 4 Chapter III: $n - n'$ and UHECR



*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 $n - n'$ and
UHECR

Introduction



Bright & Dark Sides of our Universe

- $\Omega_B \simeq 0.05$ observable matter: **electron, proton, neutron !**
- $\Omega_D \simeq 0.25$ dark matter: **WIMP? axion? sterile ν ? ...**
- $\Omega_\Lambda \simeq 0.70$ dark energy: **Λ -term? Quintessence?**
- $\Omega_R < 10^{-3}$ relativistic fraction: **relic photons and neutrinos**
- $\Omega_{\bar{B}} \approx 0$ (almost) no antimatter **only in cosmic rays/accelerators**

Matter – dark energy coincidence: $\Omega_M/\Omega_\Lambda \simeq 0.45$, ($\Omega_M = \Omega_D + \Omega_B$)
 $\rho_\Lambda \sim \text{Const.}$, $\rho_M \sim a^{-3}$; **why $\rho_M/\rho_\Lambda \sim 1$ – just Today?**

Anthropic explanation: if not *Today*, then *Yesterday* or *Tomorrow*.

Baryon and dark matter Fine Tuning: $\Omega_B/\Omega_D \simeq 0.2$
 $\rho_B \sim a^{-3}$, $\rho_D \sim a^{-3}$: **why $\rho_B/\rho_D \sim 1$ - Yesterday Today & Tomorrow?**

Baryogenesis requires BSM Physics: **(GUT-B, Lepto-B, AD-B, EW-B ...)**

Dark matter requires BSM Physics: **(Wimp, Wimpzilla, sterile ν , axion, ...)**

Different physics for B-genesis and DM?

Not very appealing: looks as Fine Tuning



Visible vs. Dark matter: $\Omega_D/\Omega_B \sim 1$?

Visible matter from Baryogenesis

B ($B - L$) & CP violation, Out-of-Equilibrium

$\rho_B = n_B m_B$, $m_B \simeq 1 \text{ GeV}$, $\eta = n_B/n_\gamma \sim 10^{-9}$

η is model dependent on several factors:

coupling constants and CP-phases, particle degrees of freedom, mass scales and out-of-equilibrium conditions, etc.



• Sakharov 1967

Dark matter: $\rho_D = n_X m_X$, but $m_X = ?$, $n_X = ?$

n_X is model dependent: DM particle mass and interaction strength (production and annihilation cross sections), freezing conditions, etc.

- Axion $m_a \sim 10^{-5} \text{ eV}$ $n_a \sim 10^4 n_\gamma$ - CDM
- Neutrinos $m_\nu \sim 10^{-1} \text{ eV}$ $n_\nu \sim n_\gamma$ - HDM (✗)
- Sterile ν' $m_{\nu'} \sim 10 \text{ keV}$ $n_{\nu'} \sim 10^{-3} n_\nu$ - WDM
- Mirror baryons $m_{B'} \simeq 1 \text{ GeV}$ $n_{B'} \sim n_B$ - ???
- WIMP $m_X \sim 1 \text{ TeV}$ $n_X \sim 10^{-3} n_B$ - CDM
- WimpZilla $m_X \sim 10^{14} \text{ GeV}$ $n_X \sim 10^{-14} n_B$ - CDM

Transforming matter and dark matter into antimatter

Zurab Berezhiani

Summary

Introduction: Dark Matter from a Parallel World

Chapter I: Neutrino - mirror neutrino mixings

Chapter II: neutron - mirror neutron mixing

Chapter III: $n - n'$ and UHECR



Dark Matter from a Parallel World

Our observable particles *very complex physics !!*

$$G = SU(3) \times SU(2) \times U(1) \quad (+ \text{SUSY ? GUT ? Seesaw ? })$$

photon, electron, nucleons (quarks), neutrinos, gluons, $W^\pm - Z$, Higgs ...

long range EM forces, confinement scale Λ_{QCD} , weak scale M_W

... matter vs. antimatter (B-L violation, CP ...)

... existence of nuclei, atoms, molecules life.... Homo Sapiens !

Best of the possible Worlds (Candid, Frank and Uncontrived)

Dark matter from parallel gauge sector ?

$$G' = SU(3)' \times SU(2)' \times U(1)' ? \quad (+ \text{SUSY ? GUT ' ? Seesaw ? })$$

photon', electron', nucleons' (quarks'), $W' - Z'$, gluons' ?

... long range EM forces, confinement at Λ'_{QCD} , weak scale M'_W ?

... asymmetric dark matter (B'-L' violation, CP ...) ?

... existence of dark nuclei, atoms, molecules ... life ... Homo Aliens ?

Another Best of the possible Worlds? (Maybe Candide had a twin?)

Call it **Yin-Yang** (in chinese, **dark-bright**) duality

describes a philosophy how opposite forces are actually complementary, interconnected and interdependent in the natural world, and how they give rise to each other as they interrelate to one another,





$$SU(3) \times SU(2) \times U(1) + SU(3)' \times SU(2)' \times U(1)'$$

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



$$G \times G'$$

Regular world

Mirror world

Elementary Particles

Elementary Particles



- Two identical gauge factors, e.g. $SU(5) \times SU(5)'$, with identical field contents and Lagrangians: $\mathcal{L}_{\text{tot}} = \mathcal{L} + \mathcal{L}' + \mathcal{L}_{\text{mix}}$
- Exact PZ_2 parity $G \leftrightarrow G'$: no new parameters in dark Lagrangian \mathcal{L}'
- MM is dark (for us) and has the same gravity
- MM is identical to standard matter, (asymmetric/dissipative/atomic) but realized in somewhat different cosmological conditions: $T'/T \ll 1$.
- New interactions between O & M particles \mathcal{L}_{mix}



– All you need is ... M world colder than ours !

For a long time M matter was not considered as a real candidate for DM: naively assuming that exactly identical microphysics of O & M worlds implies also their cosmologies are exactly identical :

- $T' = T, \quad g'_* = g_* \quad \rightarrow \quad \Delta N_\nu^{\text{eff}} = 6.15 \quad \text{vs.} \quad \Delta N_\nu^{\text{eff}} < 0.5 \quad (\text{BBN})$
- $n'_B/n'_\gamma = n_B/n_\gamma \quad (\eta' = \eta) \quad \rightarrow \quad \Omega'_B = \Omega_B \quad \text{vs.} \quad \Omega'_B/\Omega_B \simeq 5 \quad (\text{DM})$

But all is OK if : Z.B., Dolgov, Mohapatra, 1995 (*broken* PZ_2)
Z.B., Comelli, Villante, 2000 (*exact* PZ_2)

- after inflation M world was born colder than O world, $T'_R < T_R$
- any interactions between M and O particles are feeble and cannot bring two sectors into equilibrium in later epochs
- two systems evolve adiabatically (no entropy production): $T'/T \simeq \text{const}$

$T'/T < 0.5$ from BBN, but cosmological limits $T'/T < 0.2$ or so.

- $x = T'/T \ll 1 \quad \Rightarrow \quad$ in O sector 75% H + 25% ^4He
 $\Rightarrow \quad$ in M world 25% H' + 75% $^4\text{He}'$

For broken PZ_2 , DM can be compact H' atoms or n' with $m \simeq 5 \text{ GeV}$ or (sterile) mirror neutrinos $m \sim \text{few keV}$ Z.B., Dolgov, Mohapatra, 1995

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



Brief Cosmology of Mirror World

- CMB & (linear) structure formation epoch

Since $x = T'/T \ll 1$, mirror photons decouple before M-R equality:

$$z'_{\text{dec}} \simeq x^{-1} z_{\text{dec}} \simeq 1100 (T/T')$$

After that (and before M-reionization) M matter behaves as collisionless CDM and $T'/T < 0.2$ is consistent with Planck, BAO, Ly- α etc.

- **Cosmic dawn:** M world is colder (and helium dominated), the first M star can be formed earlier and reionize M sector ($z'_r \simeq 20$ or so vs $z_r = 10 \div 6$).

– EDGES 21 cm at $z \simeq 17$?

Heavy first M stars ($M \sim 10^3 M_{\odot}$) and formation of central BH – Quasars?

- **Galaxy halos?** if $\Omega'_B \simeq \Omega_B$, M matter makes ~ 20 % of DM, forming dark disk, while ~ 80 % may come from other type of CDM (WIMP?)

But perhaps 100 % ? if $\Omega'_B \simeq 5\Omega_B$: – M world is helium dominated, and the star formation and evolution can be much faster. Halos could be viewed as mirror elliptical galaxies dominated by BH and M stars, with our matter forming disks inside.

Maybe not always: Galaxies with missing DM, or too many DM, etc. ?

Because of $T' < T$, the situation $\Omega'_B \simeq 5\Omega_B$ becomes plausible in baryogenesis. So, M matter can be dark matter (as we show below)

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



CMB and LSS power spectra

Transforming matter and dark matter into antimatter

Zurab Berezhiani


Summary

Introduction: Dark Matter from a Parallel World

Chapter I: Neutrino - mirror neutrino mixings

Chapter II: neutron - mirror neutron mixing

Chapter III: $n - n'$ and UHECR





ZB, Ciarcelluti, Comelli, Villante, 2003

$$z'_{\text{dec}} \simeq x^{-1} z_{\text{dec}} \simeq 1100 (T/T')$$

← imprint of ΔN_{eff}

imprint of M-baryon oscillations



Acoustic oscillations and Silk damping scales: $x = 0.5, 0.3, 0.2$

$x < 0.2$: Galaxies with $M < 10^{8 \div 9} M_{\odot}$ will be damped




Can Mirror stars be progenitors of gravitational Wave bursts GW150914 etc. ?

Picture of Galactic halos as mirror ellipticals (Einasto density profile?), O matter disk inside (M stars = Machos)

Microlensing limits: $f \sim 20 - 40 \%$ for $M = 1 - 10 M_{\odot}$,

$f \sim 100 \%$ is allowed for $M = 20 - 200 M_{\odot}$



GW events without any optical counterpart

Massive BH compact binaries, $M \sim 10 - 100 M_{\odot}$

Can such objects be formed from MM?

M matter: 25 % Hydrogen vs 75 % Helium: M stars more compact, less opaque, less mass loses by stellar wind and evolving much faster.

Appropriate for forming such BH binaries ? And perhaps large seeds for central BH in overdense regions?



Experimental and observational manifestations

A. Cosmological implications. $T'/T < 0.2$ or so, $\Omega'_B/\Omega_B = 1 \div 5$.

Mass fraction: H' – 25%, He' – 75%, and few % of heavier C', N', O' etc.

- Mirror baryons as **asymmetric/collisional/dissipative/atomic** dark matter: M hydrogen recombination and M baryon acoustic oscillations?
- Easier formation and faster evolution of stars: Dark matter disk? Galaxy halo as mirror elliptical galaxy? Microlensing ? Neutron stars? Black Holes? Binary Black Holes? Central Black Holes?

B. Direct detection. M matter can interact with ordinary matter e.g. via kinetic mixing $\epsilon F^{\mu\nu} F'_{\mu\nu}$, etc. Mirror helium as most abundant mirror matter particles (the region of DM masses below 5 GeV is practically unexplored). Possible signals from heavier nuclei C,N,O etc.

C. Oscillation phenomena between ordinary and mirror particles.

The most interesting interaction terms in \mathcal{L}_{mix} are the ones which violate B and L of both sectors. **Neutral particles, elementary (as e.g. neutrino) or composite (as the neutron or hydrogen atom) can mix with their mass degenerate (sterile) twins:** matter disappearance (or appearance) phenomena can be observable in laboratories.

In the Early Universe, these B and/or L violating interactions can give primordial baryogenesis and dark matter genesis, with $\Omega'_B/\Omega_B = 1 \div 5$.

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



Possible portals to Mirror World: \mathcal{L}_{mix}

these terms can be limited (only) by experiment/cosmology !

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary


Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR

- **Kinetic mixing of photons** $\epsilon F^{\mu\nu} F'_{\mu\nu}$
Makes mirror matter nanocharged ($q \sim \epsilon$)
 $\epsilon < 3 \times 10^{-7}$ (EXP) $\epsilon < 5 \times 10^{-9}$ (COSM)
GUT: $\frac{1}{M^2} (\Sigma G^{\mu\nu})(\Sigma' G'_{\mu\nu})$ $\epsilon \sim \left(\frac{M_{\text{GUT}}}{M}\right)^2$



Can induce galactic magnetic fields **Z.B., Dolgov, Tkachev, 2013**

- **Neutrino-neutrino'** (active-sterile) **mixing** – discussed later
- **Neutron-neutron'** **mixing** – discussed later



Chapter I

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 $n - n'$ and
UHECR



Chapter I

Neutrino – mirror neutrino mixings




B-L violation in O and M sectors: Active-sterile mixing

- $\frac{1}{M}(l\bar{\phi})(l\bar{\phi})$ ($\Delta L = 2$) – neutrino (seesaw) masses $m_\nu \sim v^2/M$
M is the (seesaw) scale of new physics beyond EW scale.



- Neutrino -mirror neutrino mixing – (active - sterile mixing)
L and L' violation: $\frac{1}{M}(l\bar{\phi})(l\bar{\phi})$, $\frac{1}{M}(l'\bar{\phi}')(l'\bar{\phi}')$ and $\frac{1}{M}(l\bar{\phi})(l'\bar{\phi}')$



Mirror neutrinos are natural candidates for sterile neutrinos

Transforming matter and dark matter into antimatter

Zurab Berezhiani

Summary

Introduction: Dark Matter from a Parallel World

Chapter I: Neutrino - mirror neutrino mixings



Chapter II: neutron - mirror neutron mixing

Chapter III: n - n' and UHECR



Co-leptogenesis: B-L violating interactions between O and M worlds

L and L' violating operators $\frac{1}{M}(l\bar{\phi})(l\bar{\phi})$ and $\frac{1}{M}(l\bar{\phi})(l'\bar{\phi}')$ lead to processes $l\phi \rightarrow \bar{l}\bar{\phi}$ ($\Delta L = 2$) and $l\phi \rightarrow \bar{l}'\bar{\phi}'$ ($\Delta L = 1, \Delta L' = 1$)



After inflation, our world is heated and mirror world is empty: but ordinary particle scatterings transform them into mirror particles, heating also mirror world.

- These processes should be **out-of-equilibrium**
- **Violate** baryon numbers in both worlds, $B - L$ and $B' - L'$
- **Violate** also CP, given complex couplings




Green light to celebrated conditions of Sakharov



Co-leptogenesis:

Z.B. and Bento, PRL 87, 231304 (2001)

Operators $\frac{1}{M}(l\bar{\phi})(l\bar{\phi})$ and $\frac{1}{M}(l\bar{\phi})(l'\bar{\phi}')$ via seesaw mechanism – heavy RH neutrinos N_j with Majorana masses $\frac{1}{2}Mg_{jk}N_jN_k + \text{h.c.}$



Complex Yukawa couplings $Y_{ij}l_iN_j\bar{\phi} + Y'_{ij}l'_iN_j\bar{\phi}' + \text{h.c.}$

Z_2 (Xerox) symmetry $\rightarrow Y' = Y$,

PZ_2 (Mirror) symmetry $\rightarrow Y' = Y^*$

Transforming matter and dark matter into antimatter

Zurab Berezhiani

Summary

Introduction: Dark Matter from a Parallel World

Chapter I: Neutrino - mirror neutrino mixings

Chapter II: neutron - mirror neutron mixing


Chapter III: $n - n'$ and UHECR



Co-leptogenesis: Mirror Matter as Dark Anti-Matter

Z.B., arXiv:1602.08599

Hot O World \rightarrow Cold M World



$$\frac{dn_{\text{BL}}}{dt} + (3H + \Gamma)n_{\text{BL}} = \Delta\sigma n_{\text{eq}}^2$$

$$\frac{dn'_{\text{BL}}}{dt} + (3H + \Gamma')n'_{\text{BL}} = -\Delta\sigma' n_{\text{eq}}^2$$

$$\sigma(l\phi \rightarrow \bar{l}\bar{\phi}) - \sigma(\bar{l}\bar{\phi} \rightarrow l\phi) = \Delta\sigma$$

$$\sigma(l\phi \rightarrow \bar{l}'\bar{\phi}') - \sigma(\bar{l}'\bar{\phi}' \rightarrow l'\phi') = -(\Delta\sigma + \Delta\sigma')/2 \rightarrow 0 \quad (\Delta\sigma = 0)$$

$$\sigma(l\phi \rightarrow l'\phi') - \sigma(\bar{l}'\bar{\phi}' \rightarrow \bar{l}\bar{\phi}) = -(\Delta\sigma - \Delta\sigma')/2 \rightarrow \Delta\sigma \quad (0)$$

$$\Delta\sigma = \text{Im Tr}[g^{-1}(Y^\dagger Y)^* g^{-1}(Y'^\dagger Y') g^{-2}(Y^\dagger Y)] \times T^2/M^4$$

$$\Delta\sigma' = \Delta\sigma(Y \rightarrow Y')$$

$$\text{Mirror (} PZ_2 \text{): } Y' = Y^* \rightarrow \Delta\sigma' = -\Delta\sigma \rightarrow B, B' > 0$$

$$\text{Xerox (} Z_2 \text{): } Y' = Y \rightarrow \Delta\sigma' = \Delta\sigma = 0 \rightarrow B, B' = 0$$

$$\text{If } k = \left(\frac{\Gamma}{H}\right)_{T=T_R} \ll 1, \text{ neglecting } \Gamma \text{ in eqs } \rightarrow n_{\text{BL}} = n'_{\text{BL}}$$

$$\Omega'_B = \Omega_B \simeq 10^3 \frac{JM_{\text{Pl}} T_R^3}{M^4} \simeq 10^3 J \left(\frac{T_R}{10^{11} \text{ GeV}}\right)^3 \left(\frac{10^{13} \text{ GeV}}{M}\right)^4$$

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing


Chapter III:
 $n - n'$ and
UHECR



If $k = \left(\frac{\Gamma_2}{H}\right)_{T=T_R} \sim 1$, Boltzmann Eqs.

$$\frac{dn_{\text{BL}}}{dt} + (3H + \Gamma)n_{\text{BL}} = \Delta\sigma n_{\text{eq}}^2 \quad \frac{dn'_{\text{BL}}}{dt} + (3H + \Gamma')n'_{\text{BL}} = \Delta\sigma n_{\text{eq}}^2$$

should be solved with Γ :



$D(k) = \Omega_B/\Omega'_B$, $x(k) = T'/T$ for different $g_*(T_R)$ and Γ_1/Γ_2 .

So we obtain $\Omega'_B = 5\Omega_B$ when $m'_B = m_B$ but $n'_B = 5n_B$
– the reason: mirror world is colder



Chapter II

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 $n - n'$ and
UHECR

Chapter II

Neutron – mirror neutron mixing



B violating operators between O and M particles in \mathcal{L}_{mix}

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings



Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR

Ordinary quarks u, d (antiquarks \bar{u}, \bar{d})
Mirror quarks u', d' (antiquarks \bar{u}', \bar{d}')

- Neutron -mirror neutron mixing – (Active - sterile neutrons)

$$\frac{1}{M^5} (udd)(udd) \quad \& \quad \frac{1}{M^5} (udd)(u'd'd')$$



Oscillations $n \rightarrow \bar{n}$ ($\Delta B = 2$)

Oscillations $n \rightarrow \bar{n}'$ ($\Delta B = 1, \Delta B' = -1$) $B + B'$ is conserved



Neutron– antineutron mixing

Majorana mass of neutron $\epsilon(n^T C n + \bar{n}^T C \bar{n})$ violating B by two units comes from six-fermions effective operator $\frac{1}{M^5}(udd)(udd)$

Zurab Berezhiani


Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



It causes transition $n(udd) \rightarrow \bar{n}(\bar{u}\bar{d}\bar{d})$, with oscillation time $\tau = \epsilon^{-1}$

$$\epsilon = \langle n|(udd)(udd)|\bar{n} \rangle \sim \frac{\Lambda_{\text{QCD}}^6}{M^5} \sim \left(\frac{100 \text{ TeV}}{M}\right)^5 \times 10^{-25} \text{ eV}$$

Key moment: $n - \bar{n}$ oscillation destabilizes nuclei:
 $(A, Z) \rightarrow (A - 1, \bar{n}, Z) \rightarrow (A - 2, Z/Z - 1) + \pi$'s


Present bounds on ϵ from nuclear stability

$\epsilon < 1.2 \times 10^{-24} \text{ eV}$	\rightarrow	$\tau > 1.3 \times 10^8 \text{ s}$	Fe, Soudan 2002
$\epsilon < 2.5 \times 10^{-24} \text{ eV}$	\rightarrow	$\tau > 2.7 \times 10^8 \text{ s}$	O, SK 2015
$\epsilon < 7.5 \times 10^{-24} \text{ eV}$	\rightarrow	$\tau > 0.9 \times 10^8 \text{ s}$	direct limit free n



Neutron – mirror neutron mixing

Effective operator $\frac{1}{M^5}(udd)(u'd'd')$ \rightarrow mass mixing $\epsilon n C n' + \text{h.c.}$
violating B and B' – but conserving $B - B'$



$$\epsilon = \langle n | (udd)(u'd'd') | \bar{n}' \rangle \sim \frac{\Lambda_{\text{QCD}}^6}{M^5} \sim \left(\frac{1 \text{ TeV}}{M} \right)^5 \times 10^{-10} \text{ eV}$$

Key observation: $n - \bar{n}'$ oscillation cannot destabilise nuclei:
 $(A, Z) \rightarrow (A - 1, Z) + n' (p' e' \bar{\nu}')$ forbidden by energy conservation
(In principle, it can destabilise Neutron Stars)

For $m_n = m_{n'}$, $n - \bar{n}'$ oscillation can be as fast as $\epsilon^{-1} = \tau_{n\bar{n}'} \sim 1 \text{ s}$
without contradicting experimental and astrophysical limits.
(c.f. $\tau > 10 \text{ yr}$ for neutron – antineutron oscillation)

Neutron disappearance $n \rightarrow \bar{n}'$ and regeneration $n \rightarrow \bar{n}' \rightarrow n$
can be searched at small scale 'Table Top' experiments

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron – mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



Neutron – mirror neutron oscillation probability

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron – mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR


$$H = \begin{pmatrix} m_n + \mu_n \mathbf{B} \sigma & \epsilon \\ \epsilon & m_n + \mu_n \mathbf{B}' \sigma \end{pmatrix}$$

The probability of $n-n'$ transition depends on the relative orientation of magnetic and mirror-magnetic fields. The latter can exist if mirror matter is captured by the Earth

$$P_B(t) = p_B(t) + d_B(t) \cdot \cos \beta$$

$$p(t) = \frac{\sin^2 [(\omega - \omega')t]}{2\tau^2(\omega - \omega')^2} + \frac{\sin^2 [(\omega + \omega')t]}{2\tau^2(\omega + \omega')^2}$$

$$d(t) = \frac{\sin^2 [(\omega - \omega')t]}{2\tau^2(\omega - \omega')^2} - \frac{\sin^2 [(\omega + \omega')t]}{2\tau^2(\omega + \omega')^2}$$



where $\omega = \frac{1}{2}|\mu B|$ and $\omega' = \frac{1}{2}|\mu B'|$; τ - oscillation time

$$A_B^{\text{det}}(t) = \frac{N_{-B}(t) - N_B(t)}{N_{-B}(t) + N_B(t)} = N_{\text{collis}} d_B(t) \cdot \cos \beta \leftarrow \text{asymmetry}$$



A and E are expected to depend on magnetic field

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary


Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR

E.g. assume $B' = 0.12$ Gauss





Earth mirror magnetic field via the electron drag mechanism

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*



Earth can accumulate some, even tiny amount of mirror matter due to Rutherford-like scattering of mirror matter due to photon-mirror photon kinetic mixing.

Rotation of the Earth drags mirror electrons but not mirror protons (ions) since the latter are much heavier.

Circular electric currents emerge which can generate magnetic field. Modifying mirror Maxwell equations by the source (drag) term, one gets $B' \sim \epsilon^2 \times 10^{15}$ G before dynamo, and even larger after dynamo.

Such mechanism can also induce cosmological magnetic fields

Z.B., Dolgov, Tkachev, 2013



Experiments

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary


Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*

Several experiments were done, 3 by PSI group, most sensitive by the Serebrov's group at ILL, with 190 l beryllium plated trap for UCN






Experimental Strategy

To store neutrons and to measure if the amount of the survived ones depends on the magnetic field applied.

- Fill the Trap with the UCN
- Close the valve
- Wait for T_S (300 s ...)
- Open the valve
- Count the survived Neutrons



Repeat this for different orientation and values of Magnetic field.

$$N_B(T_S) = N(0) \exp [- (\Gamma + R + \bar{P}_B \nu) T_S]$$

$$\frac{N_{B1}(T_S)}{N_{B2}(T_S)} = \exp [(\bar{P}_{B2} - \bar{P}_{B1}) \nu T_S]$$

So if we find that:

$$A(B, T_S) = \frac{N_B(T_S) - N_{-B}(T_S)}{N_B(T_S) + N_{-B}(T_S)} \neq 0 \quad E(B, b, T_S) = \frac{N_B(T_S)}{N_b(T_S)} - 1 \neq 0$$



Serebrov III – Drifts of detector and monitor counts

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary


Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR


Exp. sequence: $\{B_-, B_+, B_+, B_-, B_+, B_-, B_-, B_+\}$, $B = 0.2$ G





Serebrov III – magnetic field vertical

Exp. sequence: $\{B_-, B_+, B_+, B_-, B_+, B_-, B_-, B_+\}$, $B = 0.2$ G



Analysis pointed out the presence of a signal:

$$A(B) = (7.0 \pm 1.3) \times 10^{-4} \quad \chi^2_{/dof} = 0.9 \longrightarrow 5.2\sigma$$

interpretable by $n \rightarrow n'$ with $\tau_{nn'} \sim 2 - 10s'$ and $B' \sim 0.1G$

Z.B. and Nesti, 2012



Our experiment at ILL – Z.B., Biondi, Geltenbort et al. 2018

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani


Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron – mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



4σ effect \rightarrow 2.5σ effect (conservative)



Asimov Machine: M matter can be transformed into our antimatter

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*

In normal conditions $n' \rightarrow \bar{n}$ oscillation probabilities are tiny, mirror neutrons behave nicely and do not disturb us: everyone stays on his side of the mirror



However, under well-controlled vacuum and magnetic conditions, mirror neutrons can be transformed into our antineutrons with reasonable probabilities provided that $n' \rightarrow \bar{n}$ oscillation time is indeed small ...



Now it is turn of experimentalists to turn this tale into reality ... or to exclude it – at least oscillation time $\tau_{nn'} < 10^3$ s ...
If discovered – impact can be enormous!

One could get energy out of dark matter! Two civilisations can agree to built scientific reactors/spallation facilities and exchange neutrons ... the energy produced by each reactor can turn in 1000 times more energy for parallel world .. and all live happy and healthy ...



The Gods are Subtle ... but not Malicious

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani


Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*



First Part: Against Stupidity ...

Second Part: ...The Gods Themselves ...

Third Part: ... Contend in Vain?

*"Mit der Dummheit kämpfen Götter
selbst vergebens!"* – Friedrich Schiller



Majorana Machine: transforming neutrons into antineutrons?

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR

Probability of $n \rightarrow \bar{n}$ oscillation due to direct $n - \bar{n}$ mixing (Majorana mass) is small: even in best conditions

$$P_{n\bar{n}}(t) = \left(\frac{t}{0.1 \text{ s}}\right)^2 \left(\frac{10^8 \text{ s}}{\tau_{n\bar{n}}}\right)^2 \times 10^{-18}$$

Apriori, n can have mixing with n' and \bar{n}' , and both can be fast

$$P_{nn'}(t) = \left(\frac{t}{1 \text{ s}}\right)^2 \left(\frac{1 \text{ s}}{\tau_{nn'}}\right)^2 \quad P_{n\bar{n}'}(t) = \left(\frac{t}{10 \text{ s}}\right)^2 \left(\frac{1 \text{ s}}{\tau_{n\bar{n}'}}\right)^2 \cdot 10^{-2}$$

Two effects induce $n \rightarrow \bar{n}$ transition: $n \rightarrow n' \rightarrow \bar{n}$

$$P_{n\bar{n}}(t) = P_{nn'}(t)P_{n\bar{n}'}(t) \sim 10^{-2}$$

... magnetic field must be scanned to find resonance regime!

$$m_n = 0.94 \text{ GeV} \quad E_{\text{ann}} = 2m_n c^2 = 3.3 \times 10^{-4} \text{ Erg}$$

$$\text{PSI spallation facility: } 3 \times 10^{17} \text{ n/s} \quad P \sim 10 \text{ MW}$$

ESS (European Spallation Source) would reach 1 GW – Power Plant !



Free Neutrons: Where to find Them ?

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*

Neutrons are making 1/7 fraction of baryon mass in the Universe.

But most of neutrons bound in nuclei

$n \rightarrow \bar{n}'$ or $n' \rightarrow \bar{n}$ conversions can be seen only with free neutrons.

Free neutrons are present only in

- Reactors and Spallation Facilities
- In Cosmic Rays
- During BBN epoch (fast $n' \rightarrow \bar{n}$ can solve Lithium problem)

– Transition $n \rightarrow \bar{n}'$ can take place for (gravitationally) Neutron Stars – conversion of NS into mixed ordinary/mirror NS



Chapter III

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*

Chapter III

$n - n'$ and UHECR

Z.B., Biondi, Gazizov



UHECR as protons and GZK cutoff


GZK cutoff:

Photo-pion production on the CMB if $E > E_{\text{GZK}} \approx \frac{m_{\pi} m_p}{\epsilon_{\text{CMB}}} \approx 6 \times 10^{19} \text{ eV}$:

$p + \gamma \rightarrow p + \pi^0$ (or $n + \pi^+$), $l_{\text{mfp}} \sim 5 \text{ Mpc}$ for $E > 10^{20} \text{ eV} = 100 \text{ EeV}$

Neutron decay: $n \rightarrow p + e + \bar{\nu}_e$, $l_{\text{dec}} = \left(\frac{E}{100 \text{ EeV}}\right) \text{ Mpc}$

Neutron on CMB scattering: $n + \gamma \rightarrow n + \pi^0$ (or $p + \pi^-$)



Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



UHECR as nuclei

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani


Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*







UHECR and GZK cutoff

Two giant detectors see UHECR spectra different at $E > E_{GZK}$

Pierre Auger Observatory (PAO) – South hemisphere

Telescope Array (TA) – North hemisphere

At $E < E_{GZK}$ two spectra are perfectly coincident by relative energy shift $\approx 8\%$



+ older detectors: AGASA, HiRes, etc. (all in north hemisphere)

Events with $E > 100$ EeV were observed

Cosmic Zevatrons exist in the Universe – but where is the GZK cutoff? ↻ 🔍

Transforming matter and dark matter into antimatter

Zurab Berezhiani

Summary

Introduction: Dark Matter from a Parallel World

Chapter I: Neutrino - mirror neutrino mixings

Chapter II: neutron - mirror neutron mixing

Chapter III: $n - n'$ and UHECR



But also other discrepancies are mounting ...

- **Who are carriers of UHECR ?**

PAO and TA see different chemical content: TA is compatible with protons below 10 EeV, PAO insists UHECR become heavier nuclei above $E > 10$ EeV or so – perhaps new physics ?

- **Different anistropies from North and South ?**

TA excludes isotropic distribution at $E > 57$ EeV, observes hot spot for events $E > E_{\text{GZK}}$ (which spot is colder for $E < E_{\text{GZK}}$) . PAO anisotropies not prominent: warm spot around Cen A, and small dipole for $E > 10$ EeV – are two skies really different ?

- **From where highest energy events do come ?**

$E > 100$ EeV are expected from local supercluster (Virgo, Fornax, UM, PP etc.) and closeby structures. But they do not come from these directions. TA observes small angle correlation for $E > 100$ EeV events (2 doublets), which may indicate towards strong source – from where they come?

- **Excess of cosmogenic photons ?**

Standard GZK mechanism of UHECR produces too much cascades – contradicts to Fermi-LAT photon spectrum at $E \sim 1$ TeV – local Fog ?

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings




Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



From where highest energy CR are expected ?

For protons with $E > 40, 60$ and 100 EeV (Plots by Tynyakov)



Transforming matter and dark matter into antimatter

Zurab Berezhiani

Summary

Introduction: Dark Matter from a Parallel World

Chapter I: Neutrino - mirror neutrino mixings

Chapter II: neutron - mirror neutron mixing

Chapter III: $n - n'$ and UHECR



$n - n'$ oscillation and UHECR propagation

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani


Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



Z. Berezhiani, L. Bento, *Fast neutron – Mirror neutron oscillation and ultra high energy cosmic rays*, *Phys. Lett. B* 635, 253 (2006).

- A. $p + \gamma \rightarrow p + \pi^0$ or $p + \gamma \rightarrow n + \pi^+$ $P_{pp,pn} \approx 0.5$ $l_{mfp} \sim 5$ Mpc
 B. $n \rightarrow n'$ $P_{nn'} \simeq 0.5$ $l_{osc} \sim \left(\frac{E}{100 \text{ EeV}}\right) \text{ kpc}$
 C. $n' \rightarrow p' + e' + \bar{\nu}'_e$ $l_{dec} \approx \left(\frac{E}{100 \text{ EeV}}\right) \text{ Mpc}$
 D. $p' + \gamma' \rightarrow p' + \pi'^0$ or $p' + \gamma' \rightarrow n' + \pi'^+$ $l'_{mfp} \sim (T/T')^3 l_{mfp} \gg 5$ Mpc



Ordinary and Mirror UHECR

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary


Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR

$$\frac{n'_{\text{CMB}}}{n_{\text{CMB}}} = \left(\frac{T'}{T}\right)^3 \ll 1 \quad \rightarrow \quad \frac{\ell'_{\text{mfp}}}{\ell_{\text{mfp}}} \simeq \left(\frac{T}{T'}\right)^3 \gg 1$$





$n - n'$ oscillation in the UHECR propagation

Baryon number is **not conserved** in propagation of the UHECR

$$H = \begin{pmatrix} \mu_n \mathbf{B} \sigma & \epsilon \\ \epsilon & \mu_n \mathbf{B}' \sigma \end{pmatrix} \times (\gamma = E/m_n)$$

In the intergalactic space magnetic fields are extremely small ... but for relativistic neutrons transverse component of B is enhanced by Lorentz factor: $B_{\text{tr}} = \gamma B$ ($\gamma \sim 10^{11}$ for $E \sim 100$ EeV)

Average oscillation probability:

$$P_{nn'} = \sin^2 2\theta_{nn'} \sin^2(\ell/\ell_{\text{osc}}) \simeq \frac{1}{2} [1 + Q(E)]^{-1} \quad \tan 2\theta_{nn'} = \frac{2\epsilon}{\gamma\mu_n\Delta B}$$

$$Q(E) = (\gamma\Delta B/2\epsilon)^2 \approx q \left(\frac{E}{100 \text{ EeV}}\right)^2 \quad \Delta B = |B_{\text{tr}} - B'_{\text{tr}}|$$

$$\text{If } q = 0.5 \left(\frac{\tau_{nn'}}{1 \text{ s}}\right)^2 \left(\frac{\Delta B}{1 \text{ fG}}\right)^2 < 1,$$

$n - n'$ oscillation becomes effective for $E = 100$ EeV



Are North Sky and South Sky different ?

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani



Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



The northern sky is well structured, dominated by LSC at the center, and the Great Wall and Pisces-Perseus ... The southern hemisphere is more amorphous. There is a Cetus Wall, southern part of LSC at the center, Hydra-Centaurus region but also a large and diffuse overdensity between 19 and 22 h ... "Hockey Puck" diagrams from Huchra et al., 2Mass



Swiss Cheese Model: Mirror CRs are transformed into ordinaries in nearby Voids. *Z.B., Biondi, Gazizov, 2019*

Transforming matter and dark matter into antimatter

Zurab Berezhiani

Summary


Introduction:
Dark Matter from a Parallel World

Chapter I:
Neutrino - mirror neutrino mixings

Chapter II:
neutron - mirror neutron mixing

Chapter III:
 $n - n'$ and UHECR

Adjacent Void (0–50 Mpc) $q = 0.5 \times \left(\frac{\tau_{nn'}}{1 \text{ s}}\right)^2 \left(\frac{B_{\text{tr}} - B'_{\text{tr}}}{1 \text{ fG}}\right)^2$





Swiss cheese: More distant Void (50–100 Mpc)

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani


Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR




Is northern sky (TA) is more "voidy" than the Southern sky (PAO) ?
Interestingly, some 20–30% admixture of protons above the GZK energies
improves the "chemical" fit also for PAO data [Muzio et al. 2019](#)



Arrival directions TA and PAO events of $E > 100$ EeV

TA 2008-14 ● $E > 100$ EeV, ● $79 \div 100$ EeV, ○ $57 \div 79$ EeV
PAO 2004-14 the same for $E_r = 1.1 \times E$



● 0-15 ● 15-30 ● 30-45 (d [Mpc]) ● 45-60 ● 60-75 ● 75-90

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



TA & PAO events:

correlations with sources (AGN & radiogalaxies) and mass

Transforming
matter and dark
matter into
antimatter

Zurab Berezhiani



Summary

Introduction:
Dark Matter from
a Parallel World


Chapter I:
Neutrino - mirror
neutrino mixings

Chapter II:
neutron - mirror
neutron mixing

Chapter III:
 $n - n'$ and
UHECR



Transient sources (GRB?)



$$E_r = E \text{ (TA)}, \quad E_r = 1.1 \cdot E \text{ (PAO)}$$



TA & PAO events: autocorrelations & with tracers

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani





Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 *$n - n'$ and
UHECR*





Summary

*Transforming
matter and dark
matter into
antimatter*

Zurab Berezhiani

Summary

Introduction:
Dark Matter from
a Parallel World

Chapter I:
*Neutrino - mirror
neutrino mixings*

Chapter II:
*neutron - mirror
neutron mixing*

Chapter III:
 $n - n'$ and
UHECR

The UHECR spectra observed by TA and PAO are perfectly concordant (after 10% rescaling) at energies up to 10 EeV ... but become increasingly discordant at higher energies, very strongly above the GZK cutoff (60 EeV)

The discrepancy can be due to difference between the N- and S-skies ...
N-sky is well structured, with prominent overdensities and large voids
inbetween S-sky is more amorphous with diffused galaxies ...

It is unlikely that PAO-TA discrepancy is due to different power of sources within the GZK radius (no correlation with the mass distribution at highest energies $E > 79$ EeV, no event from the Virgo or Fornax clusters, etc.)

But it can be explained in "Swiss Cheese" model: the highest energy UHECR are born from mirror UHECR in nearby holes within the GZK radius (Voids = small magnetic field) via $n' - n$ conversion

The TA signal at super-GZK energies is boosted by prominent Voids in N-hemisphere. This can also explain intermediate scale anisotropies (20-30 degrees) in the TA arrival directions while the PAO data show non ...
Interestingly, the TA/PAO spectra are concordant in the common sky ...

Our hypothesis is testable by analyzing the new data of TA/PAO at higher statistics (e.g. studying the average h_{\max} and its RMS in the common sky)