

# Update on the indication of a mass-dependent anisotropy above $10^{18.7}$ eV in the hybrid data of the Pierre Auger Observatory

*Tuesday, October 4, 2022 12:00 PM (20 minutes)*

We test for a large-scale anisotropy in the mass of arriving cosmic-ray primaries as a function of galactic latitude. The sensitivity to primary mass is obtained through the depth of shower maximum,  $X_{\max}$ , extracted from hybrid events measured over a 14-year period at the Pierre Auger Observatory. The sky is split into distinct on- and off-plane regions using the galactic latitude of each arriving cosmic ray to form two distributions of  $X_{\max}$  which are compared using an Anderson-Darling 2-samples test. A scan over roughly half of the data is used to select a lower threshold energy of  $10^{18.7}$  eV and a galactic latitude splitting at  $|b| = 30^\circ$ , which are set as a prescription for the remaining data. With these thresholds, the distribution of  $X_{\max}$  from the on-plane region is found to have a  $9.1 \pm 1.6_{-2.2}^{+2.1}$  g/cm<sup>2</sup> shallower mean and a  $5.9 \pm 2.1_{-2.5}^{+3.5}$  g/cm<sup>2</sup> narrower width than that of the off-plane region and is observed in all telescope sites independently. These differences indicate that the mean mass of primary particles arriving from the on-plane region is greater than that of those from the off-plane region. Monte Carlo studies yield a  $5.9 \times 10^{-6}$  random chance probability for the result in the independent data, lowering to a  $6.0 \times 10^{-7}$  post-penalization random chance probability when the scanned data is included. Accounting for systematic uncertainties leads to an indication for anisotropy in mass composition above  $10^{18.7}$  eV with a  $3.3\sigma$  significance. Furthermore, the result has been newly tested using additional independent FD data recovered from the quality selection process. This test disfavors the null hypothesis of the on- and off-plane regions being uniform in composition at  $2.2\sigma$  which is in good agreement with the expected sensitivity of the dataset used for this test. Possible interpretations, accompanying results and plans for further tests will be presented.

**Primary authors:** Dr MAYOTTE, Eric (Colorado School of Mines, University of Wuppertal); Dr FITOUSSI, Thomas (Karlsruhe Institute of Technology)

**Co-author:** THE PIERRE AUGER COLLABORATION

**Presenter:** Dr MAYOTTE, Eric (Colorado School of Mines, University of Wuppertal)