

MultI-messenger probe of Cosmic Ray Origins: the MICRO project

Antonio Condorelli on behalf of the MICRO project
condorelli@ijclab.in2p3.fr




Open questions

- How do burst-like signatures (GRBs, AGN-flares) fit the cosmic-ray data?
- Can current UHECR observables constrain the 3D distribution of sources?
- Do astrophysical high-energy ν , high-energy γ -rays, and UHECRs come from the same sources?




How to answer?

Public softwares to fit simultaneously UHECR observables



A diffuse flux from identical sources of UHECRs is propagated and then compared to the data at Earth: spectrum and composition.




Computation of neutrino fluxes produced along the propagation could help to constrain the parameter space.

Future steps


- Combine simulations and data to investigate if/how transient scenarios fit the observables.
- Joint accounting for composition, anisotropy, and spectral data to triangulate possible sources.
- Predict neutral fluxes from cosmic-ray propagation and interaction, including in-source propagation and comparison to current or future constraints.

Modelling the source

Hadronic interactions in UHECR simulations code




Study of hadronic interactions in CRPropa for different hadronic interaction models.




Comparison of hadronic interactions implemented in CRPropa and SimProp.

Source-propagation models



Study of UHECR productions with in-source interactions.

Multi-messenger signatures



Multi-messenger emission by a proton Flares of 10^8 GeV from blazar.

Check our website!

